Brain Research, 555 (1991) 358–359
Elsevier

Author Index

Advokat, C. and Gulati, A., Spinal transection reduces both spinal antinociception and CNS concentration of systemically administered morphine in rats, 251
Agarwal, S.K. and Calaresu, F.R., Monosynaptic connection from caudal to rostral ventrolateral medulla in the baroreceptor reflex pathway, 70
Aisenthal, L., see Le Feuvre, R.A., 245
Andrew, R.D., see Rosen, A.S., 58
Antonucci, R.F., see Giza, B.K., 1
Apkarian, A.V., see Smith, M.V., 10
Arai, A., Vanderklish, P., Kessler, M., Lee, K. and Lynch, G., A brief period of hypoxia causes proteolysis of cytoskeletal proteins in hippocampal slices, 276
Asztely, F., see Hanse, E., 173
Atrens, D.M., see Menéndez, J.A., 193
Attwell, D., see Szatkowski, M., 343

Baker, G.B., see Robertson, B., 265
Bannon, M.J., see Freeman, A.S., 281
Barbour, B., see Szatkowski, M., 343
Bazil, M.K. and Gordon, F.J., Effect of blockade of spinal NMDA receptors on sympathoexcitation and cardiovascular responses produced by cerebral ischemia, 149
Behmand, R.A., see Riachi, N.J., 19
Borkowski, J., see Mondadori, C., 107
Bridgman, P.C., see Loewy, A.D., 346
Bush, L.G., see Singh, N.A., 233

Calaresu, F.R., see Agarwal, S.K., 70
Callahan, P.M., Paris, J.M., Cunningham, K.A. and Shinnick-Gallagher, P., Decrease of GABA-immunoreactive neurons in the amygdala after electrical kindling in the rat, 335
Canonico, P.L., see Favit, A., 65
Carlen, P.L., see Niesen, C., 319
Cerruti, C., Drian, M.-J., Kamenka, J.-M. and Privat, A., Localization of dopamine carriers by BTCP, a dopamine uptake inhibitor, on nigral cells cultured in vitro, 51
Chalmers, J., see Minson, J., 326
Charlton, M.P., see Niesen, C., 319
Chattopadhyay, A.K., see Khawaja, X.Z., 164
Chiodo, L.A., see Freeman, A.S., 281
Chudler, E.H., Foote, W.E. and Poletti, C.E., Responses of cat C_1 spinal cord dorsal and ventral horn neurons to noxious and non-noxious stimulation of the head and face, 181
Connor, D.J., Langlais, P.J. and Thal, L.J., Behavioral impairments after lesions of the nucleus basalis by ibotenic acid and quisqualic acid, 84
Cornelissen, J.C.H.M., see Sijbesma, H., 295
Cotman, C.W., see Whittemore, E.R., 215
Coyle, J.T., see Vornov, J.J., 99
Cunningham, K.A., see Callahan, P.M., 335
Curtis, A.L., see Valentino, R.J., 25

Danzebrink, R.M. and Gebhart, G.F., Intrathecal coadministration of clonidine with serotonin receptor agonists produces supra-additive visceral antinociception in the rat, 35
De Guzman, C.P., Roy, R.R., Hodgson, J.A. and Edgerton, V.R., Coordination of motor pools controlling the ankle musculature in adult spinal cats during treadmill walking, 202
De Kloet, E.R., see Sijbesma, H., 295
Delrée, P., see Lefebvre, P.P., 75
Donnerer, J., Depolarization-evoked release of glutamate, aspartate and γ-aminobutyric acid from rat dorsal spinal cord slices does not originate from capsaicin-sensitive neurons, 332
Dreifuss, J.J., see Loup, F., 220
Drian, M.-J., see Cerruti, C., 51
Dubois-Dauphin, M., see Loup, F., 220
Ducret, T., see Mondadori, C., 107

Edgerton, V.R., see De Guzman, C.P., 202

Favit, A., Fiore, L., Nicoletti, F. and Canonico, P.L., Estrogen modulates stimulation of inositol phospholipid hydrolysis by norepinephrine in rat brain slices, 65
Fiore, L., see Favit, A., 65
Foote, W.E., see Chudler, E.H., 181
Frau, M., see Stefanini, E., 340
Frazer, A., see Gambarana, C., 141
Freeman, A.S., Chiodo, L.A., Lentz, S.I., Wade, K. and Bannon, M.J., Release of cholecystokinin from rat midbrain slices and modulatory effect of D_2 DA receptor stimulation, 281
Fujii, S., Saito, K., Miyakawa, H., Ito, K.I. and Kato, H., Reversal of long-term potentiation (depotentiation) induced by tetanus stimulation of the input to CA1 neurons of guinea pig hippocampal slices, 112
Fujita, S., see Usami, S.-i., 153
Fuller, L.L., see Stevens, K.E., 91

Gambarana, C., Ordway, G.A., Hauptmann, M., Tejani-Butt, S. and Frazer, A., Central administration of 1-isoproterenol in vivo induces a preferential regulation of β_2-adrenoceptors in the central nervous system of the rat, 141
Gebhart, G.F., see Danzebrink, R.M., 35
Gessa, G.L., see Stefanini, E., 340
Gibb, J.W., see Singh, N.A., 233
Giza, B.K., Scott, T.R., Sclafani, A. and Antonucci, R.F., Polysaccharides as taste stimuli: their effect in the nucleus tractus solitarius of the rat, 1
Green, I.C., see Khawaja, X.Z., 164
Green, M.A., see Li, P.P., 169
Gulati, A., see Advokat, C., 251
Gustafsson, B., see Hanse, E., 173

Hanse, E., Asztely, F., Wigström, H. and Gustafsson, B., Long-lasting potentiations evoked by a brief heterosynaptic tetanus in the guinea pig dentate gyrus in vitro, 173
Hanson, G.R., see Singh, N.A., 233
Harik, S.I., see Riachi, N.J., 19
Hauptmann, M., see Gambarana, C., 141
Hodgson, J.A., see De Guzman, C.P., 202
Höhmann, C.F., see Jucker, M., 305
Hozawa, J., see Usami, S.-i., 153

Ingram, D.K., see Jucker, M., 305
Ishida-Yamamoto, A., see Shida, T., 288
Ito, K.-i., see Fujii, S., 112

Jin, H., see Usami, S.-i., 153
Jucker, M., Kleinman, H.K., Höhmann, C.F., Ordy, J.M. and Ingram, D.K., Distinct immunoreactivity to 110 kDa laminin-binding protein in adult and lesioned rat forebrain, 305

Kalant, H., see Mihic, S.J., 259
Kamenka, J.-M., see Cerruti, C., 51
Kaneko, T., see Minson, J., 326
Kapoor, V., see Minson, J., 326
Kato, H., see Fujii, S., 112
Kelly, P.A.T., see McBean, D.E., 159
Kessler, M., see Arai, A., 276
Khawaja, X.Z., Chattopadhyay, A.K. and Green, I.C., Increased β-endorphin and dynorphin concentrations in discrete hypothalamic regions of genetically obese (*ob/ob*) mice, 164
Kleinman, H.K., see Jucker, M., 305

Langlais, P.J., see Connor, D.J., 84
Lee, K., see Arai, A., 276
Lefebvre, P.P., Weber, T., Leprince, P., Rigo, J.-M., Delrée, P., Rogister, B. and Moonen, G., Kainate and NMDA toxicity for cultured developing and adult rat spiral ganglion neurons: further evidence for a glutamatergic excitatory neurotransmission at the inner hair cell synapse, 75
Le Feuvre, R.A., Aisenthal, L. and Rothwell, N.J., Involvement of corticotrophin releasing factor (CRF) in the thermogenic and anorexic actions of serotonin (5-HT) and related compounds, 245
Lentz, S.I., see Freeman, A.S., 281
Leprince, P., see Lefebvre, P.P., 75
Leung, L.S. and Shen, B., Hippocampal CA1 evoked response and radial 8-arm

Brain Research, 555 (1991) 358–359
Elsevier

Author Index

Advokat, C. and Gulati, A., Spinal transection reduces both spinal antinociception and CNS concentration of systemically administered morphine in rats, 251
Agarwal, S.K. and Calaresu, F.R., Monosynaptic connection from caudal to rostral ventrolateral medulla in the baroreceptor reflex pathway, 70
Aisenthal, L., see Le Feuvre, R.A., 245
Andrew, R.D., see Rosen, A.S., 58
Antonucci, R.F., see Giza, B.K., 1
Apkarian, A.V., see Smith, M.V., 10
Arai, A., Vanderklish, P., Kessler, M., Lee, K. and Lynch, G., A brief period of hypoxia causes proteolysis of cytoskeletal proteins in hippocampal slices, 276
Asztely, F., see Hanse, E., 173
Atrens, D.M., see Menéndez, J.A., 193
Attwell, D., see Szatkowski, M., 343

Baker, G.B., see Robertson, B., 265
Bannon, M.J., see Freeman, A.S., 281
Barbour, B., see Szatkowski, M., 343
Bazil, M.K. and Gordon, F.J., Effect of blockade of spinal NMDA receptors on sympathoexcitation and cardiovascular responses produced by cerebral ischemia, 149
Behmand, R.A., see Riachi, N.J., 19
Borkowski, J., see Mondadori, C., 107
Bridgman, P.C., see Loewy, A.D., 346
Bush, L.G., see Singh, N.A., 233

Calaresu, F.R., see Agarwal, S.K., 70
Callahan, P.M., Paris, J.M., Cunningham, K.A. and Shinnick-Gallagher, P., Decrease of GABA-immunoreactive neurons in the amygdala after electrical kindling in the rat, 335
Canonico, P.L., see Favit, A., 65
Carlen, P.L., see Niesen, C., 319
Cerruti, C., Drian, M.-J., Kamenka, J.-M. and Privat, A., Localization of dopamine carriers by BTCP, a dopamine uptake inhibitor, on nigral cells cultured in vitro, 51
Chalmers, J., see Minson, J., 326
Charlton, M.P., see Niesen, C., 319
Chattopadhyay, A.K., see Khawaja, X.Z., 164
Chiodo, L.A., see Freeman, A.S., 281
Chudler, E.H., Foote, W.E. and Poletti, C.E., Responses of cat C_1 spinal cord dorsal and ventral horn neurons to noxious and non-noxious stimulation of the head and face, 181
Connor, D.J., Langlais, P.J. and Thal, L.J., Behavioral impairments after lesions of the nucleus basalis by ibotenic acid and quisqualic acid, 84
Cornelissen, J.C.H.M., see Sijbesma, H., 295
Cotman, C.W., see Whittemore, E.R., 215
Coyle, J.T., see Vornov, J.J., 99
Cunningham, K.A., see Callahan, P.M., 335
Curtis, A.L., see Valentino, R.J., 25

Danzebrink, R.M. and Gebhart, G.F., Intrathecal coadministration of clonidine with serotonin receptor agonists produces supra-additive visceral antinociception in the rat, 35
De Guzman, C.P., Roy, R.R., Hodgson, J.A. and Edgerton, V.R., Coordination of motor pools controlling the ankle musculature in adult spinal cats during treadmill walking, 202
De Kloet, E.R., see Sijbesma, H., 295
Delrée, P., see Lefebvre, P.P., 75
Donnerer, J., Depolarization-evoked release of glutamate, aspartate and γ-aminobutyric acid from rat dorsal spinal cord slices does not originate from capsaicin-sensitive neurons, 332
Dreifuss, J.J., see Loup, F., 220
Drian, M.-J., see Cerruti, C., 51
Dubois-Dauphin, M., see Loup, F., 220
Ducret, T., see Mondadori, C., 107

Edgerton, V.R., see De Guzman, C.P., 202

Favit, A., Fiore, L., Nicoletti, F. and Canonico, P.L., Estrogen modulates stimulation of inositol phospholipid hydrolysis by norepinephrine in rat brain slices, 65
Fiore, L., see Favit, A., 65
Foote, W.E., see Chudler, E.H., 181
Frau, M., see Stefanini, E., 340
Frazer, A., see Gambarana, C., 141
Freeman, A.S., Chiodo, L.A., Lentz, S.I., Wade, K. and Bannon, M.J., Release of cholecystokinin from rat midbrain slices and modulatory effect of D_2 DA receptor stimulation, 281
Fujii, S., Saito, K., Miyakawa, H., Ito, K.I. and Kato, H., Reversal of long-term potentiation (depotentiation) induced by tetanus stimulation of the input to CA1 neurons of guinea pig hippocampal slices, 112
Fujita, S., see Usami, S.-i., 153
Fuller, L.L., see Stevens, K.E., 91

Gambarana, C., Ordway, G.A., Hauptmann, M., Tejani-Butt, S. and Frazer, A., Central administration of 1-isoproterenol in vivo induces a preferential regulation of β_2-adrenoceptors in the central nervous system of the rat, 141
Gebhart, G.F., see Danzebrink, R.M., 35
Gessa, G.L., see Stefanini, E., 340
Gibb, J.W., see Singh, N.A., 233
Giza, B.K., Scott, T.R., Sclafani, A. and Antonucci, R.F., Polysaccharides as taste stimuli: their effect in the nucleus tractus solitarius of the rat, 1
Green, I.C., see Khawaja, X.Z., 164
Green, M.A., see Li, P.P., 169
Gulati, A., see Advokat, C., 251
Gustafsson, B., see Hanse, E., 173

Hanse, E., Asztely, F., Wigström, H. and Gustafsson, B., Long-lasting potentiations evoked by a brief heterosynaptic tetanus in the guinea pig dentate gyrus in vitro, 173
Hanson, G.R., see Singh, N.A., 233
Harik, S.I., see Riachi, N.J., 19
Hauptmann, M., see Gambarana, C., 141
Hodgson, J.A., see De Guzman, C.P., 202
Höhmann, C.F., see Jucker, M., 305
Hozawa, J., see Usami, S.-i., 153

Ingram, D.K., see Jucker, M., 305
Ishida-Yamamoto, A., see Shida, T., 288
Ito, K.-i., see Fujii, S., 112

Jin, H., see Usami, S.-i., 153
Jucker, M., Kleinman, H.K., Höhmann, C.F., Ordy, J.M. and Ingram, D.K., Distinct immunoreactivity to 110 kDa laminin-binding protein in adult and lesioned rat forebrain, 305

Kalant, H., see Mihic, S.J., 259
Kamenka, J.-M., see Cerruti, C., 51
Kaneko, T., see Minson, J., 326
Kapoor, V., see Minson, J., 326
Kato, H., see Fujii, S., 112
Kelly, P.A.T., see McBean, D.E., 159
Kessler, M., see Arai, A., 276
Khawaja, X.Z., Chattopadhyay, A.K. and Green, I.C., Increased β-endorphin and dynorphin concentrations in discrete hypothalamic regions of genetically obese (*ob/ob*) mice, 164
Kleinman, H.K., see Jucker, M., 305

Langlais, P.J., see Connor, D.J., 84
Lee, K., see Arai, A., 276
Lefebvre, P.P., Weber, T., Leprince, P., Rigo, J.-M., Delrée, P., Rogister, B. and Moonen, G., Kainate and NMDA toxicity for cultured developing and adult rat spiral ganglion neurons: further evidence for a glutamatergic excitatory neurotransmission at the inner hair cell synapse, 75
Le Feuvre, R.A., Aisenthal, L. and Rothwell, N.J., Involvement of corticotrophin releasing factor (CRF) in the thermogenic and anorexic actions of serotonin (5-HT) and related compounds, 245
Lentz, S.I., see Freeman, A.S., 281
Leprince, P., see Lefebvre, P.P., 75
Leung, L.S. and Shen, B., Hippocampal CA1 evoked response and radial 8-arm

maze performance after hippocampal kindling, 353
Li, P.P., Green, M.A. and Warsh, J.J., A study of cerebellar inositol 1,4,5-trisphosphate receptor following climbing and parallel fibre deafferentation, 169
Llewellyn-Smith, I., see Minson, J., 326
Loewy, A.D., Bridgman, P.C. and Mettenleiter, T.C., β-galactosidase expressing recombinant pseudorabies virus for light and electron microscopic study of transneuronally labeled CNS neurons, 346
Lookingland, K.J., see Tian, Y., 135
Loup, F., Tribollet, E., Dubois-Dauphin, M. and Dreifuss, J.J., Localization of high-affinity binding sites for oxytocin and vasopressin in the human brain. An autoradiographic study., 220
Lynch, G., see Arai, A., 276

Martini, M., see Rossi, M.L., 123
Matsubara, A., see Usami, S.-i., 153
McBean, D.E., Sharkey, J., Ritchie, I.M. and Kelly, P.A.T., Cerebrovascular and functional consequences of 5-HT_{1A} receptor activation, 159
Menéndez, J.A. and Atrens, D.M., Insulin and the paraventricular hypothalamus: modulation of energy balance, 193
Mettenleiter, T.C., see Loewy, A.D., 346
Midgley, L.P., see Singh, N.A., 233
Mihic, S.J., Wu, P.H. and Kalant, H., Differences among effects of sedative-hypnotic drugs on GABA-mediated chloride flux: quench flow studies, 259
Minson, J., Pilowsky, P., Llewellyn-Smith, I., Kaneko, T., Kapoor, V. and Chalmers, J., Glutamate in spinally projecting neurons of the rostral ventral medulla, 326
Miyakawa, H., see Fujii, S., 112
Mondadori, C., Ducret, T. and Borkowski, J., How long does 'memory consolidation' take? New compounds can improve retention performance, even if administered up to 24 hours after the learning experience, 107
Moonen, G., see Lefebvre, P.P., 75
Moore, K.E., see Tian, Y., 135

Nicoletti, F., see Favit, A., 65
Niesen, C., Charlton, M.P. and Carlen, P.L., Postsynaptic and presynaptic effects of the calcium chelator BAPTA on synaptic transmission in rat hippocampal dentate granule neurons, 319
Noebels, J.L., see Qiao, X., 43

Ordway, G.A., see Gambarana, C., 141
Ordy, J.M., see Jucker, M., 305
Otani, H., see Yoshioka, T., 177

Page, M.E., see Valentino, R.J., 25
Pai, K.S. and Ravindranath, V., Protection and potentiation of MPTP-induced toxicity by cytochrome P-450 inhibitors and inducer: in vitro studies with brain slices, 239
Paris, J.M., see Callahan, P.M., 335
Pilowsky, P., see Minson, J., 326
Poletti, C.E., see Chudler, E.H., 181
Privat, A., see Cerruti, C., 51

Qiao, X. and Noebels, J.L., Genetic and phenotypic heterogeneity of inherited spike-wave epilepsy: two mutant gene loci with independent cerebral excitability defects, 43

Ravindranath, V., see Pai, K.S., 239
Riachi, N.J., Behmand, R.A. and Harik, S.I., Correlation of MPTP neurotoxicity in vivo with oxidation of MPTP by the brain and blood–brain barrier in vitro in five rat strains, 19
Rigo, J.-M., see Lefebvre, P.P., 75
Ritchie, I.M., see McBean, D.E., 159
Robertson, B., Baker, G.B. and Vanderwolf, C.H., The effects of serotonergic stimulation on hippocampal and neocortical slow waves and behavior, 265
Rogister, B., see Lefebvre, P.P., 75
Rose, G.M., see Stevens, K.E., 91
Rosen, A.S. and Andrew, R.D., Glucose concentration inversely alters neocortical slice excitability through an osmotic effect, 58
Rossi, M.L. and Martini, M., Efferent control of posterior canal afferent receptor discharge in the frog labyrinth, 123
Rothwell, N.J., see Le Feuvre, R.A., 245
Roy, R.R., see De Guzman, C.P., 202

Saito, K., see Fujii, S., 112
Schipper, J., see Sijbesma, H., 295
Sclafani, A., see Giza, B.K., 1
Scott, T.R., see Giza, B.K., 1
Senba, E., see Shida, T., 288
Sharkey, J., see McBean, D.E., 159
Shen, B., see Leung, L.S., 353
Shida, T., Ueda, Y., Ishida-Yamamoto, A., Senba, E. and Tohyama, M., Enkephalinergic sympathetic and parasympathetic innervation of the rat submandibular and sublingual glands, 288
Shinnick-Gallagher, P., see Callahan, P.M., 335
Shinohara, H., see Yoshioka, T., 177
Sijbesma, H., Schipper, J., Cornelissen, J.C.H.M. and De Kloet, E.R., Species differences in the distribution of central 5-HT_1 binding sites: a comparative autoradiographic study between rat and guinea pig, 295
Singh, N.A., Midgley, L.P., Bush, L.G., Gibb, J.W. and Hanson, G.R., *N*-Methyl-D-aspartate receptors mediate dopamine-induced changes in extrapyramidal and limbic dynorphin systems, 233
Smith, M.V. and Apkarian, A.V., Thalamically projecting cells of the lateral cervical nucleus in monkey, 10
Stefanini, E., Frau, M. and Gessa, G.L., Increase in D_2 dopamine receptors in the substantia nigra after chronic (–)-sulpiride treatment, 340
Stevens, K.E., Fuller, L.L. and Rose, G.M., Dopaminergic and noradrenergic modulation of amphetamine-induced changes in auditory gating, 91
Straznicky, C., see Wye-Dvorak, J., 313
Szatkowski, M., Barbour, B. and Attwell, D., The potassium-dependence of excitatory amino acid transport: resolution of a paradox, 343

Tanaka, O., see Yoshioka, T., 177
Tazawa, M., see Usami, S.-i., 153
Tejani-Butt, S., see Gambarana, C., 141
Thal, L.J., see Connor, D.J., 84
Tian, Y., Lookingland, K.J. and Moore, K.E., Contribution of noradrenergic neurons to 3,4-dihydroxyphenylacetic acid concentrations in the regions of the rat brain containing incertohypothalamic dopaminergic neurons, 135
Tohyama, M., see Shida, T., 288
Tribollet, E., see Loup, F., 220

Ueda, Y., see Shida, T., 288
Usami, S.-i., Hozawa, J., Tazawa, M., Jin, H., Matsubara, A. and Fujita, S., Localization of substance P-like immunoreactivity in guinea pig vestibular endorgans and the vestibular ganglion, 153

Valentino, R.J., Page, M.E. and Curtis, A.L., Activation of noradrenergic locus coeruleus neurons by hemodynamic stress is due to local release of corticotropin-releasing factor, 25
Vanderklish, P., see Arai, A., 276
Vanderwolf, C.H., see Robertson, B., 265
Vornov, J.J. and Coyle, J.T., Enhancement of NMDA receptor-mediated neurotoxicity in the hippocampal slice by depolarization and ischemia, 99

Wade, K., see Freeman, A.S., 281
Warsh, J.J., see Li, P.P., 169
Weber, T., see Lefebvre, P.P., 75
Whittemore, E.R. and Cotman, C.W., Agonists selective for phosphoinositide-coupled receptors sensitize neurons to depolarization by L-2-amino-4-phosphonobutanoic acid (L-AP4), 215
Wigström, H., see Hanse, E., 173
Wu, P.H., see Mihic, S.J., 259
Wye-Dvorak, J. and Straznicky, C., Retinal distribution of ganglion cells which project to the ipsilateral optic tectum in *Bufo marinus*, 313

Yoshioka, T., Tanaka, O., Otani, H. and Shinohara, H., Histochemically demonstrable phosphotyrosine protein phosphatase in the rat hippocampal formation, 177